

# Anti-Human NECTIN4/PVRL4 Antibody (SAA0404)

## Summary


---

<b>Catalog No.</b>	FHJ41010
<b>Clone ID</b>	SAA0404
<b>Host species</b>	Human
<b>Conjugation</b>	Unconjugated
<b>Species reactivity</b>	Human
<b>Form</b>	Liquid
<b>Storage buffer</b>	0.01M PBS, pH 7.4, 0.09% Sodium azide.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	>95% as determined by SDS-PAGE.
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG1, kappa
<b>Applications</b>	FCM
<b>Target</b>	LNIR, NECTIN4, PRR4 receptor-related protein 4, Nectin cell adhesion molecule 4, Ig superfamily receptor LNIR, Nectin-4, PVRL4
<b>Purification</b>	Protein A/G purified from cell culture supernatant.
<b>Endotoxin level</b>	Please contact with the lab for this information.
<b>Accession</b>	Q96NY8
<b>Stability and Storage</b>	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. Store at 4°C short term (1-2 weeks). Store at -20°C 12 months. Store at -80°C long term.


**Note**

For research use only.

## Data Image



Flow-cytometry



Flow-cytometry

Flow-cytometry using anti-human NECTIN4 antibody. NECTIN4 Transfected CHO cells were stained with an irrelevant antibody (Blue Histogram) or an anti-human NECTIN4 antibody monoclonal antibody (Catalog # FHJ41010, Green Histogram) at a concentration of 5 µg/ml for 30 mins at RT. After washing, bound antibody was detected using a FITC conjugated goat anti-human antibody (Catalog # PHB96441) and cells analysed on a NovoCyt Flow Cytometer.

Flow-cytometry using anti-human NECTIN4 antibody. MCF7 cells were stained with an irrelevant antibody (Blue Histogram) or an anti-human NECTIN4 antibody monoclonal antibody (Catalog # FHJ41010, Green Histogram) at a concentration of 5 µg/ml for 30 mins at RT. After washing, bound antibody was detected using a FITC conjugated goat anti-human antibody (Catalog # PHB96441) and cells analysed on a NovoCyt Flow Cytometer.