

Anti-VEGFA Polyclonal Antibody

Summary

Catalog No. PHD12601

Host species Rabbit

Tested applications ELISA: 1:4000-1:8000, IHC: 1:50-1:100, WB: 1:1000-1:4000

Species reactivity Human, Mouse, Rat

Immunogen E. coli - derived recombinant Human VEGFA (Ala27-Arg232).

Form Liquid

Storage buffer 0.01M PBS, pH 7.4, 50% Glycerol, 0.05% Proclin 300.

Clonality Polyclonal

Isotype IgG

Applications ELISA, IHC, WB

Vascular endothelial growth factor A, VPF, VEGFA, VEGF, Vascular **Target**

permeability factor, VEGF-A

Purification Purified by antigen affinity column.

Accession P15692

Use a manual defrost freezer and avoid repeated freeze thaw cycles. Store


Stability and Storage at 2 to 8°C for frequent use. Store at -20 to -80°C for twelve months from

the date of receipt.


Note For research use only.

Data Image

Recombinant Proteins & Antibodies



WESTERN BLOT



Immunocytochemistry/ Immunofluorescence

Recombinant Protein lysates were subjected to SDS PAGE followed by western blot with VEGFA antibody (PHD12601) at 1 µg/ml.

Lane 1: Recombinant protein

Second Ab: Goat Anti-Rabbit IgG H&L Polyclonal antibody, HRP (PTB96431) at 0.1 µg/mL.

Predict MW: 25 kDa Observed MW: 25 kDa

VEGFA in HepG2 Cell Line.

The HepG2 cells were fixed with 4% paraformaldehyde (20 min), and then blocked with 5% goat serum for 1h. And the cells were incubated for 2h at 37°C with VEGFA (PHD12601) at 8 μg/ml. The section was then incubated with Goat Anti-Rabbit IgG (Alexa Fluor-488) preabsorbed at 1/100 dilution (Shown in green) for 1 hour at room temperature. Nuclear DNA was labelled with DAPI (shown in blue).