

## Anti-CD281/TLR1 Polyclonal Antibody

## **Summary**

Catalog No. PHH08901

**Host species** Rabbit

**Tested applications** ELISA: 1:4000-1:8000, IHC: 1:50-1:100, WB: 1:1000-1:4000

**Species reactivity** Human

**Immunogen** E. coli - derived recombinant Human CD281/TLR1 (Asn625-Lys786).

**Form** Liquid

Storage buffer 0.01M PBS, pH 7.4, 50% Glycerol, 0.05% Proclin 300.

**Clonality** Polyclonal

**Isotype** IgG

**Applications** ELISA, IHC, WB

Toll/interleukin-1 receptor-like protein,CD281,Toll-like receptor **Target** 

1,TIL,KIAA0012,TLR1

**Purification** Purified by antigen affinity column.

Accession Q15399


Use a manual defrost freezer and avoid repeated freeze thaw cycles. Store

Stability and Storage at 2 to 8°C for frequent use. Store at -20 to -80°C for twelve months from


the date of receipt.

Note For research use only.

## Data Image



WESTERN BLOT



Immunocytochemistry/ Immunofluorescence

Recombinant Protein lysates were subjected to SDS PAGE followed by western blot with CD281/TLR1 antibody (PHH08901) at 1 µg/ml.

Lane 1: Recombinant protein

Second Ab: Goat Anti-Rabbit IgG H&L Polyclonal antibody, HRP (PTB96431) at 0.1 µg/mL.

Predict MW: 21 kDa Observed MW: 21 kDa

CD281 / TLR1 in Hela Cell Line.

The Hela cells were fixed with 4% paraformaldehyde (20 min), and then blocked with 5% goat serum for 1h. And the cells were incubated for 2h at 37°C with CD281 / TLR1 (PHH08901) at 2.6 µg/ml. The section was then incubated with Goat Anti-Rabbit IgG (Alexa Fluor-594) preabsorbed at 1/100 dilution (Shown in red) for 1 hour at room temperature. Nuclear DNA was labelled with DAPI (shown in blue).