

# Anti-SLC9A3R1 Antibody (R1D25)

## Summary

---


<b>Catalog No.</b>	RHA28601
<b>Clone ID</b>	R1D25
<b>Host species</b>	Rabbit
<b>Tested applications</b>	IHC: 1:100-1:200, WB: 1:1000-1:2000
<b>Species reactivity</b>	Human, Rat
<b>Form</b>	Liquid
<b>Storage buffer</b>	0.01M PBS, pH 7.4, 0.05% BSA, 50% Glycerol, 0.05% Sodium azide.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	>95% by SDS-PAGE.
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG
<b>Applications</b>	IHC, WB
<b>Target</b>	NHERF1, Sodium-hydrogen exchanger regulatory factor 1, SLC9A3R1, Solute carrier family 9 isoform A3 regulatory factor 1, NHERF, EBP50, Ezrin-radixin-moesin-binding phosphoprotein 50, Na(+)/H(+) exchange regulatory cofactor NHE-RF1, NHERF-1, Regulatory cofactor of Na(+)/H(+) exchanger
<b>Purification</b>	Protein A/G purified from cell culture supernatant.
<b>Endotoxin level</b>	Please contact with the lab for this information.
<b>Accession</b>	O14745

**Stability and Storage**

Use a manual defrost freezer and avoid repeated freeze thaw cycles. Store at 4 °C for frequent use. Store at -20 °C for twelve months from the date of receipt.


**Note**

For research use only.

**Data Image**


Western blot

Western blot analysis of EBP50 expression in (1) HepG2 cell lysate; (2) PC12 cell lysate.



Western blot

All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



Western blot

All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.