

Anti-CYP17A1 Antibody (R1J95)

Summary


Catalog No.	RHC12902
Clone ID	R1J95
Host species	Rabbit
Tested applications	FCM: 1:20-1:100, IF: 1:50-1:200, IP: 1:20-1:50, WB: 1:1000-1:2000
Species reactivity	Human, Mouse, Rat
Form	Liquid
Storage buffer	0.01M PBS, pH 7.4, 0.05% BSA, 50% Glycerol, 0.05% Sodium azide.
Concentration	1 mg/ml
Purity	>95% by SDS-PAGE.
Clonality	Monoclonal
Isotype	IgG
Applications	FCM, IF, IP, WB
Target	Steroid 17-alpha-hydroxylase/17,20 lyase, CYPXVII, Cytochrome P450 17A1, Cytochrome P450-C17, Steroid 17-alpha-monooxygenase, 17-alpha-hydroxyprogesterone aldolase, CYP17A1, S17AH, Cytochrome P450c17, CYP17
Purification	Protein A/G purified from cell culture supernatant.
Endotoxin level	Please contact with the lab for this information.
Accession	P05093

Stability and Storage

Use a manual defrost freezer and avoid repeated freeze thaw cycles. Store at 4 °C for frequent use. Store at -20 °C for twelve months from the date of receipt.


Note

For research use only.

Data Image


Western blot

Western blot analysis of Cytochrome P450 17A1 expression in Jurkat cell lysate.




Western blot

All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.




Western blot

All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.




Western blot

All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.




Western blot

All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



Western blot

All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



Western blot

All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.