

# Anti-RIOX2 Antibody (R2M66)

## Summary

---


<b>Catalog No.</b>	RHK63701
<b>Clone ID</b>	R2M66
<b>Host species</b>	Rabbit
<b>Tested applications</b>	IF: 1:50-1:200, WB: 1:1000-1:2000
<b>Species reactivity</b>	Human, Mouse
<b>Form</b>	Liquid
<b>Storage buffer</b>	0.01M PBS, pH 7.4, 0.05% BSA, 50% Glycerol, 0.05% Sodium azide.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	>95% by SDS-PAGE.
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG
<b>Applications</b>	IF, WB
<b>Target</b>	ROX, MDIG, Nucleolar protein 52, Bifunctional lysine-specific demethylase and histidyl-hydroxylase MINA, Ribosomal oxygenase 2, MINA, Histone lysine demethylase MINA, Ribosomal oxygenase MINA, MYC-induced nuclear antigen, RIOX2, 60S ribosomal protein L27a histidine hydroxylase, Mineral dust-induced gene protein, NO52, MINA53
<b>Purification</b>	Protein A/G purified from cell culture supernatant.
<b>Endotoxin level</b>	Please contact with the lab for this information.
<b>Accession</b>	Q8IUF8

**Stability and Storage**


Use a manual defrost freezer and avoid repeated freeze thaw cycles. Store at 4 °C for frequent use. Store at -20 °C for twelve months from the date of receipt.

**Note**


For research use only.

**Data Image**

All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



Western blot

Western blot analysis of MINA53 expression in A431 cell lysate.