

Anti-EFHD1 Antibody (R4D66)


Summary

Catalog No.	RHP12201
Clone ID	R4D66
Host species	Mouse
Tested applications	IF: 1:50-1:200, IHC: 1:50-1:100, WB: 1:500-1:1000
Species reactivity	Human, Mouse, Rat
Form	Liquid
Storage buffer	0.01M PBS, pH 7.4, 0.5% BSA, 0.05% Sodium Azide and 50% Glycerol.
Concentration	1 mg/ml
Purity	>95% as determined by SDS-PAGE.
Clonality	Monoclonal
Isotype	IgG1
Applications	IF, IHC, WB
Target	EF-hand domain-containing protein 1, SWS2, EF-hand domain-containing protein D1, EFHD1, Swiprosin-2
Purification	Protein A/G purified from cell culture supernatant.
Endotoxin level	Please contact with the lab for this information.
Accession	Q9BUP0
Stability and Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. Store at 4°C short term (1-2 weeks). Store at -20°C 12 months. Store at -80°C long term.

Note

For research use only.

Data Image




Immunofluorescence

Immunofluorescence analysis of EFHD1 in mouse lung using EFHD1 antibody(3G2)(red),and DAPI (blue).




Immunofluorescence

Immunofluorescence analysis of EFHD1 in Hela using EFHD1 antibody(Left) and DAPI (Right).




Immunohistochemical

Immunohistochemistry analysis of paraffin-embedded Human uterus tissue using EFHD1 antibody.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.Negative control was used by secondary antibody only.




Immunohistochemical

Immunohistochemistry analysis of paraffin-embedded Human tonsils using EFHD1 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Negative control was used by secondary antibody only.




Immunohistochemical

Immunohistochemistry analysis of paraffin-embedded mouse testis tissue using EFHD1 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Negative control was used by secondary antibody only.



Immunohistochemical

Immunohistochemistry analysis of paraffin-embedded mouse testis tissue using EFHD1 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval. Negative control was used by secondary antibody only.



Western blot

Western blot analysis of EFHD1 in mouse spleen, rat spleen tissue lysates using EFHD1 antibody.