


# Anti-ARNTL Polyclonal Antibody

## Summary

---

<b>Catalog No.</b>	PHA09301
<b>Host species</b>	Rabbit
<b>Tested applications</b>	ELISA: 1:4000-1:8000, IHC: 1:50-1:100, WB: 1:1000-1:4000
<b>Species reactivity</b>	Human, Mouse, Rat
<b>Immunogen</b>	E. coli - derived recombinant Human ARNTL (Glu48-Leu320).
<b>Form</b>	Liquid
<b>Storage buffer</b>	0.01M PBS, pH 7.4, 50% Glycerol, 0.05% Proclin 300.
<b>Clonality</b>	Polyclonal
<b>Isotype</b>	IgG
<b>Applications</b>	ELISA, IHC, WB
<b>Target</b>	Basic-helix-loop-helix-PAS protein MOP3,ARNTL,Member of PAS protein 3,PAS domain-containing protein 3,BMAL1,Aryl hydrocarbon receptor nuclear translocator-like protein 1,BHLHE5,Class E basic helix-loop-helix protein 5,bHLHe5,MOP3,Brain and muscle ARNT-like 1,bHLH-PAS protein JAP3,PASD3
<b>Purification</b>	Purified by antigen affinity column.
<b>Accession</b>	O00327
<b>Stability and Storage</b>	Use a manual defrost freezer and avoid repeated freeze thaw cycles. Store at 2 to 8°C for frequent use. Store at -20 to -80°C for twelve months from the date of receipt.
<b>Note</b>	For research use only.

## Data Image



WESTERN BLOT

Recombinant Protein lysates were subjected to SDS PAGE followed by western blot with ARNTL antibody (PHA09301) at 1 µg/ml.

Lane 1: Recombinant protein


Second Ab: Goat Anti-Rabbit IgG H&L Polyclonal antibody, HRP (PTB96431) at 0.1 µg/mL.

Predict MW: 33 kDa


Observed MW: 33 kDa

ARNTL in NIH3T3 Cell Line.

The NIH3T3 cells were fixed with 4% paraformaldehyde (20 min), and then blocked with 5% goat serum for 1h. And the cells were incubated for 2h at 37°C with ARNTL (PHA09301) at 13.2 µg/ml. The section was then incubated with Goat Anti-Rabbit IgG (Alexa Fluor-488) preabsorbed at 1/100 dilution (Shown in green) for 1 hour at room temperature. Nuclear DNA was labelled with DAPI (shown in blue).



Immunocytochemistry/ Immunofluorescence



Immunocytochemistry/ Immunofluorescence

**ARNTL in NIH3T3 Cell Line.**

The NIH3T3 cells were fixed with 4% paraformaldehyde (20 min), and then blocked with 5% goat serum for 1h. And the cells were incubated for 2h at 37°C with ARNTL (PHA09301) at 13.2 µg/ml. The section was then incubated with Goat Anti-Rabbit IgG (Alexa Fluor-488) preabsorbed at 1/100 dilution (Shown in green) for 1 hour at room temperature. Nuclear DNA was labelled with DAPI (shown in blue).