

Anti-Human CD112R/PVRIG Antibody (CPA.7.021)

Summary

Catalog No.	FHK16210
Clone ID	CPA.7.021
Host species	Mouse
-	
Conjugation	Unconjugated
Species reactivity	Human
Form	Liquid
Storage buffer	0.01M PBS, pH 7.4.
Concentration	1 mg/ml
Purity	>95% as determined by SDS-PAGE.
Clonality	Monoclonal
Isotype	IgG1, kappa
Applications	FCM
Target	C7orf15, CD112R, CD112 receptor receptor-related immunoglobulin domain-containing protein, PVRIG, Transmembrane protein PVRIG
Purification	Protein A/G purified from cell culture supernatant.
Endotoxin level	Please contact with the lab for this information.
Accession	Q6DKI7
Stability and Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. Store at 4°C short term (1-2 weeks). Store at -20°C 12 months. Store at - 80°C long term.




🍸 AntibodySystem


Recombinant Proteins & Antibodies

Note

For research use only.

Data Image





Flow-cytometry using anti-human CD112R antibody.Jurkat cells were stained with an irrelevant antibody (Blue Histogram) or an anti-human CD112R antibody monoclonal antibody (Catalog # FHK16210 ,Green Histogram) at a concentration of 5 μ g/ml for 30 mins at RT. After washing, bound antibody was detected using a FITC conjugated goat anti-mouse antibody (Catalog # PMB96441) and cells analysed on a NovoCyte Flow Cytometer.


Flow-cytometry using anti-human CD112R antibody.MOLT-4 cells were stained with an irrelevant antibody (Blue Histogram) or an antihuman CD112R antibody monoclonal antibody (Catalog # FHK16210 ,Green Histogram) at a concentration of 5 µg/ml for 30 mins at RT. After washing, bound antibody was detected using a FITC conjugated goat anti-mouse antibody (Catalog # PMB96441) and cells analysed on a NovoCyte Flow Cytometer.

 $\mathbf{\nabla}$





Recombinant Proteins & Antibodies



Flow-cytometry using anti-human CD112R antibody.Raji cells were stained with an irrelevant antibody (Blue Histogram) or an anti-human CD112R antibody monoclonal antibody (Catalog # FHK16210 ,Green Histogram) at a concentration of 5 µg/ml for 30 mins at RT. After washing, bound antibody was detected using a FITC conjugated goat anti-mouse antibody (Catalog # PMB96441) and cells analysed on a NovoCyte Flow Cytometer.



 $\mathbf{\nabla}$

