


Anti-Human CD220/INSR Antibody (AGT-181)

Summary


| | |
|------------------------------|---|
| Catalog No. | FHC19110 |
| Clone ID | AGT-181 |
| Host species | Mouse |
| Conjugation | Unconjugated |
| Species reactivity | Human |
| Form | Liquid |
| Storage buffer | 0.01M PBS, pH 7.4. |
| Concentration | 1 mg/ml |
| Purity | >95% as determined by SDS-PAGE. |
| Clonality | Monoclonal |
| Isotype | IgG2a, kappa |
| Applications | FCM |
| Target | INSR, IR, CD220, Insulin receptor |
| Purification | Protein A/G purified from cell culture supernatant. |
| Endotoxin level | Please contact with the lab for this information. |
| Accession | P06213 |
| Stability and Storage | Use a manual defrost freezer and avoid repeated freeze-thaw cycles. Store at 4°C short term (1-2 weeks). Store at -20°C 12 months. Store at -80°C long term. |
| Note | For research use only. |

Data Image



Flow-cytometry

Flow-cytometry using anti-human CD220 antibody. 293T cells were stained with an irrelevant antibody (Blue Histogram) or an anti-human CD220 antibody monoclonal antibody (Catalog # FHC19110, Green Histogram) at a concentration of 5 µg/ml for 30 mins at RT. After washing, bound antibody was detected using a FITC conjugated goat anti-mouse antibody (Catalog # PMB96441) and cells analysed on a NovoCyt Flow Cytometer.



Flow-cytometry

Flow-cytometry using anti-human CD220 antibody. MCF7 cells were stained with an irrelevant antibody (Blue Histogram) or an anti-human CD220 antibody monoclonal antibody (Catalog # FHC19110, Green Histogram) at a concentration of 5 µg/ml for 30 mins at RT. After washing, bound antibody was detected using a FITC conjugated goat anti-mouse antibody (Catalog # PMB96441) and cells analysed on a NovoCyt Flow Cytometer.