

# Anti-Human CD276 Antibody (Mab 8H9)

## Summary

---

<b>Catalog No.</b>	FHJ04020
<b>Clone ID</b>	Mab 8H9
<b>Host species</b>	Mouse
<b>Conjugation</b>	Unconjugated
<b>Species reactivity</b>	Human
<b>Form</b>	Liquid
<b>Storage buffer</b>	0.01M PBS, pH 7.4.
<b>Concentration</b>	3.19 mg/ml
<b>Purity</b>	>95% as determined by SDS-PAGE.
<b>Clonality</b>	Monoclonal
<b>Isotype</b>	IgG1, kappa
<b>Applications</b>	FCM
<b>Target</b>	CD276, B7-H3, Costimulatory molecule, B7H3, 4Ig-B7-H3, B7 homolog 3, CD276 antigen
<b>Purification</b>	Protein A/G purified from cell culture supernatant.
<b>Endotoxin level</b>	Please contact with the lab for this information.
<b>Accession</b>	Q5ZPR3
<b>Stability and Storage</b>	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. Store at 4°C short term (1-2 weeks). Store at -20°C 12 months. Store at -80°C long term.


**Note**

For research use only.

For research use only


## Data Image

---




### Flow-cytometry

Flow-cytometry using anti-human CD276 antibody.293T cells were stained with an irrelevant antibody (Blue Histogram) or an anti-human CD276 antibody monoclonal antibody (Catalog # FHJ04020 ,Green Histogram) at a concentration of 5  $\mu$ g/ml for 30 mins at RT. After washing, bound antibody was detected using a FITC conjugated goat anti-mouse antibody (Catalog # PMB96441) and cells analysed on a NovoCyte Flow Cytometer.



### Flow-cytometry

Flow-cytometry using anti-human CD276 antibody.PC-3 cells were stained with an irrelevant antibody (Blue Histogram) or an anti-human CD276 antibody monoclonal antibody (Catalog # FHJ04020 ,Green Histogram) at a concentration of 5  $\mu$ g/ml for 30 mins at RT. After washing, bound antibody was detected using a FITC conjugated goat anti-mouse antibody (Catalog # PMB96441) and cells analysed on a NovoCyte Flow Cytometer.



### Bioactivity

Detects CD276 Protein in indirect ELISA.